

COMPUTABILITY, GÖDEL'S INCOMPLETENESS THEOREM, AND AN INHERENT LIMIT ON THE PREDICTABILITY OF EVOLUTION

Troy Day^{1,2}

1. Department of Mathematics and Statistics,
Jeffery Hall, Queen's University,
Kingston, ON, K7L 3N6, Canada
tday@mast.queensu.ca, Ph: 613-533-2431, Fax: 613-533-2964

2. Department of Biology, Queen's University
Kingston, ON, K7L 3N6, Canada

October 14, 2011

Abstract

The process of evolutionary diversification unfolds in a vast genotypic space of potential outcomes. During the past century there have been remarkable advances in the development of theory for this diversification (Fisher, 1930; Wright, 1984; Hofbauer and Sigmund, 1988; Lynch and Walsh, 1998; Bürger, 2000; Ewens, 2004; Barton et al., 2007), and the theory's success rests, in part, on the scope of its applicability. A great deal of this theory focuses on a relatively small subset of the space of potential genotypes, chosen largely based on historical or contemporary patterns, and then predicts the evolutionary dynamics within this pre-defined set. To what extent can such an approach be pushed to a broader perspective that accounts for the potential open-endedness of evolutionary diversification? There have been a number of significant theoretical developments along these lines (Gillespie, 1984; Fontana and Buss, 1994; Szathmary, 1995; Maynard Smith and Szathmáry, 1995; Wagner and Altenberg, 1996; Orr, 1998; Stadler et al., 2001; Yedid and Bell, 2002; Orr, 2002; Wagner and Stadler, 2003; Fernando and Rowe, 2007; Nowak and Ohtsuki, 2008; Joyce et al., 2008; Ohtsuki and Nowak, 2009; Manapat et al., 2009) but the question of how far such theory can be pushed has not been addressed. Here a theorem is proven demonstrating that, because of the digital nature of inheritance, there are inherent limits on the kinds of questions that can be answered using such an approach. In particular, even in extremely simple evolutionary systems a complete theory accounting for the potential open-endedness of evolution is unattainable unless evolution is progressive. The theorem is closely related to Gödel's Incompleteness Theorem (Gödel, 1931; Nagel and Newman, 1958; Davis, 1965; van Heijenoort ed., 1967) and to the Halting Problem from computability theory (Turing, 1936; Cutland, 1980).

1 Introduction

2 Much of evolutionary theory is, in an important sense, fundamentally historical. The
3 process of evolutionary diversification unfolds in a vast genotypic space of potential
4 outcomes, and explores some parts of this space and not others. Nevertheless, a great deal
5 of current theory restricts attention to a relatively small subset of this space, chosen largely
6 based on historical or contemporary patterns, and then predicts evolutionary dynamics.
7 Although this can work well for making short-term predictions, ultimately it must fail once
8 evolution gives rise to genuinely novel genotypes lying outside this predefined set
9 (Yedid and Bell, 2002).

10 This potential limitation on the predictive ability of many models of evolution has been
11 noted on various occasions throughout the development of evolutionary theory (Levinton,
12 1988; Fontana and Buss, 1994; Wagner and Altenberg, 1996; Yedid and Bell, 2002),
13 perhaps most famously by Dutch biologist Hugo DeVries when he remarked that “Natural
14 selection may explain the survival of the fittest, but it cannot explain the arrival of the
15 fittest” (DeVries, 1904). Such statements hint at the notion that many models of evolution
16 are what we might call ‘local’, or ‘closed’, in the sense that they focus attention on a very
17 small (local) region of the evolutionary tree and do not account for the possibility that
18 evolution is an open-ended process.

19 The distinction between ‘closed’ and ‘open-ended’ models of evolution will be discussed in
20 more detail below, but in recent years there have been several interesting studies published
21 that are beginning to push the boundaries of analyses towards what we might naturally
22 call open-ended models. These studies include models of abstract replicator populations
23 (Fontana and Buss, 1994; Szathmary, 1995; Nowak and Ohtsuki, 2008;
24 Ohtsuki and Nowak, 2009; Manapat et al., 2009), models exploring the space of
25 evolutionary possibilities (Fontana and Schuster, 1998b; Stadler et al., 2001;

Wagner and Stadler, 2003), analyses of evolutionary transitions (Maynard Smith and Szathmáry, 1995; Fontana and Schuster, 1998a), models for predicting the distribution of allelic effects during evolution (Gillespie, 1984; Orr, 1998, 2002; Joyce et al., 2008), and studies of evolvability (Wagner and Altenberg, 1996). Similarly, there have also been many *in silico* and artificial life experiments that explore generic, emergent, properties of evolution (Fontana and Buss, 1994; Lenski et al., 1999; Yedid and Bell, 2001; Wilke et al., 2001; Yedid and Bell, 2002; Lenski et al., 2003; Chow et al., 2004; Ostrowski et al., 2007; Fernando and Rowe, 2007; Yedid et al., 2008, 2009). In general these analyses have demonstrated that, once we allow for more open-ended evolution, a much richer suite of evolutionary possibilities arises.

The above studies collectively suggest that accounting for open-ended evolution in theory can yield interesting new insights, and it can also yield new testable predictions (Gillespie, 1984; Orr, 1998, 2002; Joyce et al., 2008). Nevertheless, there is still a relative paucity of theoretical studies that allow for open-ended evolution, and so we might expect that much is yet to be learned by broadening evolutionary theory further in this way. My purpose with this article is therefore twofold. First, I simply wish to highlight the fact that there is an important distinction to be made between open-ended versus closed models of evolution (defined more precisely below), and to suggest that open-ended models might more faithfully represent the evolutionary process. Second, and more significantly, I wish to consider whether a push towards a predictive theory that embraces the potential open-endedness of evolution is likely to face additional obstacles, over and above those faced by closed models of evolution. Put another way, I ask the question: To what extent is the development of a predictive, open-ended evolutionary theory possible?

Although a complete answer to the above question is not possible, in what follows I will provide at least a partial answer. Furthermore, I demonstrate that this answer has interesting connections to the Halting Problem from computability theory and to Gödel's

Incompleteness Theorem from mathematical logic. In particular, I will use results from these areas to prove a theorem that formally links the concept of progressive evolution to the possibility of developing such a predictive open-ended theory. There remains debate over if, and when, evolution might be progressive (Dawkins, 1997; Gould, 1997; Adami et al., 2000) and part of this debate stems from the lack of a precise yet general definition of progression. Thus, another way to view the results presented here is as providing such a definition. I will return to this point more fully in the discussion.

A Motivating Example

To sharpen the focus on these somewhat abstract ideas, it is worth beginning with a concrete motivating example involving evolutionary prediction. This section does so, focussing primarily on the broad conceptual issues involved. The section that follows then addresses these issues more precisely.

Consider trying to use evolutionary theory to predict the dynamics of human influenza. Specifically, consider trying to answer the following question: is it likely that a pandemic with the 1918 Spanish influenza strain will ever occur again? This is obviously a difficult, and still somewhat loosely defined, question so let's narrow things down further. One reason we might be skeptical about our ability to make such predictions is because of uncertainty in initial conditions and parameter values, as well as uncertainty about the evolutionary processes involved. In other words, perhaps we lack all of the information required to make such predictions. Furthermore, unexpected contingencies might thwart what would otherwise be accurate predictions. For example, an unanticipated volcanic eruption might temporarily alter commercial air travel patterns, and this might thereby alter the epidemiological and evolutionary dynamics of influenza.

These practical limitations are clearly important, but are they the only obstacle to making

76 accurate evolutionary predictions or are there other, ‘inherent’, limitations as well. Does
77 the difficulty of making evolutionary predictions stem simply from our lack of knowledge of
78 the evolutionary processes involved or are there reasons why, even in principle, such
79 evolutionary predictions are not possible?

80 It is this latter question that is the focus of this article, and therefore I will, at least
81 temporarily, put the above practical concerns aside. Specifically, let’s assume that we can
82 build a model that adequately captures all of the relevant evolutionary processes, and that
83 we can obtain all parameter estimates necessary to use such a model. Without getting too
84 much into the specifics, one of the first things we would need to decide is the relevant strain
85 space for the model. The simplest scenario would consider only two strains (e.g., the 1918
86 strain and the current, predominant, strain). More sophisticated scenarios might instead
87 include several strains that are thought to be important in the dynamics. In either case,
88 both such resulting models would be ‘closed’ in the sense described in the introduction
89 because they focus only on a finite (and relatively small) number of strains. Furthermore,
90 given that there is a discrete and finite number of people who can be infected at any given
91 time, there is then also a finite (and relatively small) number of possible evolutionary
92 outcomes. As will be detailed more precisely later, this then implies that the process will
93 either reach a steady state or it will display periodic behaviour (see Appendix 5). Hence, if
94 a closed model is an accurate description of the evolutionary process, then in principle we
95 can answer the above question by simply running the model until one of these two outcomes
96 occurs. At that point we need only observe whether or not a 1918 Spanish flu pandemic
97 ever occurred during the run of the model (or if it occurred with significant probability).

98 But what if the evolutionary process is, instead, open-ended? To explore this possibility we
99 need to be more specific about what is meant by open-ended. Consider again the influenza
100 example. Influenza A has a genome size of more than 12,000 nucleotides, and therefore the
101 number of possible genotypes is enormous. To gain some perspective on just how many

genotypes are possible, let's restrict attention to only the smallest of the eight genomic segments of influenza. In this case there are then only approximately 800 nucleotides and therefore approximately 4^{800} different possible genotypes. To put this number in perspective, it is approximately 10^{400} times larger than the estimated number of atoms in the universe. For a model to be open-ended it would have to allow for such a vast set of possible evolutionary outcomes so that, as in reality, evolutionary change could continue unabated, producing potentially novel outcomes essentially indefinitely. The simplest way we might try to capture this theroretically is to assume that the space of possible genotypes is infinite.

Given these considerations, if evolutionary theory is to capture an open-ended evolutionary process, then its state space must be effectively infinite. This is necessary but it is not a sufficient condition for open-ended evolution. For example, many stochastic Markovian models in population genetics have an infinite state space (e.g., the infinite alleles model; Kimura and Crow (1964)) but nevertheless do not display open-ended evolution. Rather, further assumptions are often made, such as the assumption that the Markov chain is irreducible and positively recurrent. These assumptions are usually made primarily for mathematical convenience but they rule out the possibility of open-ended evolution since they then guarantee the existence a single unique equilibrium or stationary distribution. As a result, such models cannot capture the possibility that evolutionary change might continue indefinitely.

What if we relax these assumptions and allow for truly open-ended evolution in the theory that we develop? Are there then even further problems associated with making evolutionary predictions? For example, does this make answering the question about influenza evolution laid out at the start of this section more difficult? You might suspect that the answer is 'yes'; at least, the approach suggested above for closed models will no longer suffice because the evolutionary process is no longer guaranteed to settle down to an

equilibrium or stationary distribution. Thus, the best we can possibly hope for is that there is some way to prove, using the structure of the model, whether or not such an outcome will occur. Thus, all practical difficulties of predicting evolution aside, it is not obvious whether we can answer the above sort of question about influenza evolution, even in principle.

These issues are now starting to tread heavily into the fields of computability and mathematical logic and, roughly speaking, a theory that can answer the above kind of question about influenza evolution is referred to as a negation-complete theory. This terminology reflects the idea that the theory is complete in the sense of one being able to determine whether a given statement is true, or whether its formal negation is true instead. For example, in the context of influenza, a negation-complete theory would be able to predict whether the statement ‘the Spanish flu will happen again’ is true or whether its formal negation ‘it is not true that the Spanish flu will happen again’ is true instead. More generally, a negation-complete evolutionary theory would be one from which we could determine those parts of genotypic space will be explored by evolution and those that will not.

Is such a negation-complete theory possible once we allow for open-ended evolution? In the remainder of this article I show that the answer to this question is closely related to the idea of progressive evolution. In particular, even if the system of evolution were simple enough for us to understand everything about how its genetic composition changes from one generation to the next, the following theorem is proven:

Theorem: A negation-complete evolutionary theory is possible if, and only if, the evolutionary process is progressive.

The above theorem will be made more precise shortly, but as already alluded to above, it stems from the fact that DNA affords evolution a mechanism of digital inheritance. As Maynard Smith and Szathmáry have noted (Maynard Smith and Szathmáry, 1995) the

combinatorial complexity that arises thereby allows evolution to be effectively open-ended. Indeed, as will be argued below, digital inheritance allows one to characterize evolution (i.e., the change in genetic composition of a population) as a dynamical system on the natural numbers, and therefore the theorem proved below holds for any such dynamical system, not just those meant to model evolution. As a result, the theorem is closely related to other results from mathematics and computer science; namely Gödel’s Incompleteness Theorem (Gödel, 1931; Nagel and Newman, 1958; Davis, 1965; van Heijenoort ed., 1967) and to the Halting Problem from computability theory (Turing, 1936; Cutland, 1980).

Statement and Proof of Theorem

In order to give precision to the above theorem, we must specify what is meant by ‘the evolutionary process’, as well as what it means for evolutionary theory to be negation-complete. The goal is to determine if, even in extremely simple evolutionary processes, there is some inherent limitation on evolutionary theory.

To this end, consider a simplified evolutionary process in which there is a well-mixed population of replicators with some maximal population size, and in which each replicator contains a single piece of DNA. This genetic code can mutate in both composition, and in length, with no pre-imposed bounds. Suppose that each replicator survives and reproduces in a way that depends only on the current genetic composition of the population. For additional simplicity, suppose that generations are discrete. All conclusions hold if events occur in continuous time instead (Appendix 5). Finally, for simplicity of exposition, I will usually assume that the evolutionary dynamics are deterministic in the main text. Again, all results generalize to the case of stochastic evolutionary dynamics, albeit with a few additional assumptions (Appendix 5).

With the above evolutionary dynamic, the genetic composition of the system will evolve

177 over time, and we can characterize the state of the system at any time by the number of
178 each type of replicator (e.g., the number of infections with each possible genotype of
179 influenza). The goal then is to determine if it is possible to construct an evolutionary
180 theory that can predict which parts of the space of potential evolutionary outcomes will be
181 explored during evolutionary diversification, and which will not. Formally, the results
182 presented below are valid for any theory whose derived statements are recursively
183 enumerable. Axiomatic theories are one such example but (roughly speaking) any
184 theoretical approach that can, in principle, be implemented by a computer falls into this
185 category (Appendix 1). Indeed, the statement and proof of the theorem relies on several
186 ideas from computability theory (Appendix 2).

187 The digital nature of inheritance provided by DNA means that, in principle, the number of
188 distinct kinds of replicators that are possible is discrete and unbounded, a property
189 Maynard Smith and Szathmáry refer to as ‘indefinite’ heredity
190 (Maynard Smith and Szathmáry, 1995). It is indefinite heredity that allows for open-ended
191 evolution. As a result, in principle, the set of possible population states during evolution is
192 isomorphic to the positive integers; i.e., there exists a one-to-one correspondence between
193 the set of possible population states and the positive integers. Such sets are called
194 denumerable, and in fact the set of population states is effectively denumerable in a
195 computability sense (Appendix 3). Thus we can effectively assign a unique integer-valued
196 ‘code’ to every possible population state.

197 In practice, of course, there are limits on the number of kinds of replicators possible, if only
198 because of a finite pool of the required chemical building blocks. Nevertheless, as
199 mentioned earlier the combinatorial nature of indefinite heredity means that the actual
200 number of possible population states is so large as to be effectively infinite. For simplicity
201 of exposition, it is assumed in the main text that the set of possible population states is
202 truly infinite; however, Appendix 6 makes the notion of ‘effectively infinite’ precise and

provides the analogous results for this case.

With the above coding we can formalize evolution mathematically as a mapping of the positive integers to themselves. For example, in the deterministic case we might start with a model (e.g., a mapping F) that tells us the number of individuals of each genotype in the next time step, as a function of the current numbers. Then, under the above coding, if $E(n)$ denotes the population state (formally, its integer code number) at time n , the model can be recast as a single-variable, integer, mapping $E(n + 1) = G(E(n))$ for some function G , along with some initial condition. Similarly, in the stochastic case, if we start with a probabilistic mapping F , then it can be recast as a mapping $E(n + 1) = H(E(n))$ where H gives the probability distribution over the set of code numbers in the next time step as a function of its current distribution (and E is then a vector of probabilities over the integers). Therefore, in general, we can view the evolutionary trajectory as being simply an integer-valued function with an integer-valued argument. Of course, different ways of coding the population states will correspond to different maps, G or H , and thus different functions $E(n)$. Also note that the domain of G or H need not be all of the positive integers, and in fact different initial conditions might give rise to different domains as well. This would correspond to there being different basins of attraction in the evolutionary process.

It is also worth noting that, although we have assumed the evolutionary mapping (i.e., G or H) is a function of the current genetic composition of the population only, we can relax this assumption and allow evolutionary change to depend on other aspects of the environment as well. In particular, we might expand our definition of ‘population state’ to include both genetic state, and the state of other variables associated with the environment in which the genes exist. Again, as long as such generalized processes can be recast as dynamical systems on the natural numbers, all of the results presented here continue to hold.

The above arguments illustrate how we can view evolution as a dynamical system on the

229 natural numbers, and they also now allow us to formalize the notion of open-ended
 230 evolution. In the deterministic setting evolution is open-ended if the mapping G never
 231 revisits a previously visited state. Likewise, in the stochastic setting, evolution is
 232 open-ended if the mapping H always admits at least one new state each generation with
 233 positive probability.

234 Because we can view evolution as a dynamical system on the natural numbers,
 235 evolutionary theory can be viewed as a set of specific rules for manipulating and deducing
 236 statements about such numbers. Computability theory deals with functions that map
 237 positive integers to themselves, and thus provides a natural set of tools to analyze the
 238 problem. A function is called ‘computable’ if there exists some algorithmic procedure that
 239 can be followed to evaluate the function in a finite number of steps (Appendix 2).

240 Again, focusing on the deterministic case, given the assumption that we are able to predict
 241 the state of the population from one time step to the next, the function $E(n)$ is
 242 computable (see Appendix 2). Furthermore, the set of all computable functions is
 243 denumerable (Cutland, 1980). Therefore, denoting the k^{th} such function by $\phi_k(n)$, it is
 244 clear the evolutionary process, $E(n)$, must correspond to a member of this set. Denote this
 245 specific member by $\phi_E(n)$, and again note that, if we change the integer-coding used to
 246 identify specific population states, we will obtain a different function $\hat{E}(n)$, and thus a
 247 different member of the set, $\phi_{\hat{E}}(n)$ (Fig. 1).

248 During evolution, a set of population states will be visited over time (in the stochastic case
 249 we consider a state as being visited if the probability of it occurring at some point is larger
 250 than a threshold value; Appendix 5). These will be referred to as ‘evolutionarily attainable’
 251 states. In terms of our formalism, this corresponds to the function $\phi_E(n)$ taking on various
 252 values of its range, R_E , as n increases (Fig. 1). A negation-complete evolutionary theory
 253 would be one that can determine whether a code, x , satisfies $x \in R_E$ or whether it satisfies
 254 $x \notin R_E$ instead. In the language of computability theory, this corresponds to asking

255 whether the predicate ' $x \in R_E$ ' is decidable (Appendix 2; (Cutland, 1980)). In terms of the
 256 influenza example presented earlier, if x is the population state corresponding to a
 257 pandemic with the 1918 strain, then the statement 'the Spanish flu will happen again'
 258 corresponds to the number-theoretic statement $x \in R_E$. Likewise, the statement 'it is not
 259 true that the Spanish flu will happen again' corresponds to the number-theoretic statement
 260 $x \notin R_E$.

261 Lastly, we can give a precise definition of progressive evolution. Intuitively, evolution is
 262 progressive if there is some quantifiable characteristic of the population that increases
 263 through evolutionary time. In terms of the above formalization, this means there is a way
 264 to recode the population states such that the code number increases during evolution.
 265 Formally, evolution is progressive if there exists a computable, one-to-one, coding of the
 266 population states by positive integers, $\hat{C}$, such that the corresponding description of the
 267 evolutionary process, $\phi_{\hat{E}}(n)$, satisfies $\phi_{\hat{E}}(n+1) > \phi_{\hat{E}}(n)$ for all n . Again, in terms of the
 268 influenza example presented earlier, if evolution were progressive, then there would be some
 269 way to *a priori* code the population states such that, as influenza evolution occurs, the
 270 code number of the population increases (I will return to this definition of progression in
 271 more detail in the discussion).

272 We can now rephrase the theorem in terms of precise, technical, language:

273 *Theorem: ' $x \in R_E$ ' is decidable if, and only if, there exists a computable, one-to-one,*
 274 *coding of the population states by positive integers, $\hat{C}$, such that the corresponding*
 275 *description of the evolutionary process, $\phi_{\hat{E}}(n)$, satisfies $\phi_{\hat{E}}(n+1) > \phi_{\hat{E}}(n)$ for all n .*

276 Proof (Figure 1; see Appendices 2 and 4 for additional details):

277 Part 1: If there exists a coding $\hat{C}$ such that $\phi_{\hat{E}}(n+1) > \phi_{\hat{E}}(n)$ for all n then the predicate
 278 ' $x \in R_E$ ' is decidable.

279 By hypothesis there exists a computable bijection $\hat{C}$ such that, for the corresponding
 280 description of the evolutionary process, $\phi_{\hat{E}}(n+1) > \phi_{\hat{E}}(n)$ for all n . For any population
 281 state, x , in the original coding, let $\hat{x}$ be the corresponding code under the bijection $\hat{C}$, and
 282 define $z(\hat{x}) = \mu i(\phi_{\hat{E}}(i) \geq \hat{x})$, where $\mu i(H(i))$ denotes the minimum value of i for which the
 283 argument $H(i)$ is true (Appendix 2). Further, define $R_k(n) = \{x : \phi_k(i) = x, i \leq n\}$ (i.e.,
 284 the range of $\phi_k(n)$ visited by step n ; Appendix 2). Clearly ' $\hat{x} \in R_{\hat{E}}(z(\hat{x}))$ ' is decidable since
 285 $R_{\hat{E}}(z(\hat{x}))$ is finite and can be enumerated, and furthermore $\hat{x} \in R_{\hat{E}}(z(\hat{x})) \Leftrightarrow \hat{x} \in R_{\hat{E}}$ owing
 286 to the progressive nature of evolution. Therefore, ' $\hat{x} \in R_{\hat{E}}$ ' is decidable as well. Finally,
 287 using S denote the set of population states that are evolutionarily attainable, we have that
 288 $\hat{x} \in R_{\hat{E}} \Leftrightarrow \hat{C}^{-1}\hat{x} \in S \Leftrightarrow C\hat{C}^{-1}\hat{x} \in R_E$. Noting that, by definition, $x = C\hat{C}^{-1}\hat{x}$, we obtain
 289 $\hat{x} \in R_{\hat{E}} \Leftrightarrow x \in R_E$. Thus, ' $x \in R_E$ ' is decidable as well.

290 Part 2: If the predicate ' $x \in R_E$ ' is decidable then there exists a coding $\hat{C}$ such that
 291 $\phi_{\hat{E}}(n+1) > \phi_{\hat{E}}(n)$ for all n .

292 We can construct the required computable bijection between population states and an
 293 appropriate coding as follows. First, take any effective coding of population states. By
 294 hypothesis ' $x \in R_E$ ' is decidable and therefore we can proceed through the population
 295 states, x , in increasing order, applying the following algorithm:

296 (i) if $x \notin R_E$ and it is the k^{th} such state up to that point, use the k^{th} odd number as its
 297 new code.

298 (ii) if $x \in R_E$, calculate $\mu i(\phi_E(i) = x)$, and use the i^{th} even number as its new code.

299 Thus, $R_{\hat{E}}$ is the set of even numbers, and they are visited in increasing order as evolution
 300 proceeds. In particular, using $\hat{C}C^{-1}$ to denote the above mapping described in points (i)
 301 and (ii), where C^{-1} is the inverse mapping of the coding that generated x (i.e., it takes
 302 code x and returns the corresponding population state, s), we have

303 $\phi_{\hat{E}}(n+1) = \hat{C}C^{-1}\phi_E(n+1) = 2(n+1)$. The last equality follows from the fact that

304 $\hat{C}C^{-1}\phi_E(n+1)$ determines the time at which state $\phi_E(n+1)$ occurs (which is $n+1$), and
 305 assigns it a new code equal to twice this value (point (ii) above). Therefore
 306 $\phi_{\hat{E}}(n+1) > \phi_{\hat{E}}(n) \forall n$.
 307 Q.E.D.

308 Discussion

309 This article has two main goals. The first goal is to highlight the distinction between
 310 open-ended versus closed models of evolution, and to suggest that open-ended models
 311 might better capture real evolutionary processes. The second goal is to explore the extent
 312 to which the development of a predictive, open-ended theory of evolution is possible. The
 313 above theorem illustrates that there is an interesting connection between this question and
 314 analyses from computability theory and mathematical logic. It also draws a formal
 315 connection between the extent to which such a theory is possible and the notion of
 316 progressive evolution.

317 Because the theorem states an equivalence relationship between the possibility of
 318 developing a negation-complete theory and progressive evolution, it can be read in two
 319 distinct ways. First, it states that if evolution is progressive then a negation-complete
 320 theory is possible. This is, perhaps, not too surprising. If evolution is progressive then
 321 there would be a good deal of regularity to the process that one ought to be able to exploit
 322 in constructing theory. The second way to read the theorem is from the perspective of the
 323 reverse implication. This is somewhat more surprising; it states that if evolution is not
 324 progressive then a negation-complete theory will not be possible.

325 These results rest on the fact that digital inheritance allows evolution to be open-ended
 326 (Maynard Smith and Szathmary, 1995). If, instead, the hereditary system allowed for only

a finite number of discrete possible types, then evolution would either display periodic behaviour or would reach an equilibrium (possibly with stochastic fluctuations; Appendix 5). A negation-complete theory of evolution would then be trivially possible in such cases because, in principle, we could simply develop a finite list of all evolutionary outcomes that can occur (as described in the influenza example earlier).

Of course, despite the existence of digital inheritance, there is nevertheless presumably a bound on the number of population states possible for a variety of reasons. Even so, however, the combinatorial nature of digital inheritance means that the number of possible population states might be considered effectively infinite. An analogous theorem can be proven in such cases by replacing the notion of infinite with a precise notion of effectively infinite instead (Appendix 6). Likewise, although the main results of the text assume that evolution is deterministic, an analogous theorem holds that accounts for the inherently stochastic nature of the evolutionary process (Appendix 5).

The notion of progressive evolution is somewhat slippery, and there does not exist a general yet precise definition of progression that is universally agreed upon. As a result, this has led to disagreement over the extent to which progressive evolution occurs (Dawkins, 1997; Gould, 1997). A complete discussion of the idea of progressive evolution is beyond the scope of this article but a few points are worth making here.

Most discussions of progressive evolution involve quantities like mean fitness, body size, complexity, or other relatively conspicuous biological measurements. Many such discussions also are retrospective in the sense that they look at historical patterns when attempting to find patterns of progression. But both of these aspects of discussions of progression are problematic. First, although it would be nice to readily identify some obvious, and biologically meaningful, characteristic of a population that changes in a directional way, there is no reason to expect that we have currently thought of all the possibilities. Thus, when defining progression, it would seem desirable to do so in a very general way, leaving

open the possibility that some biologically interesting, but as yet undiscovered quantity increases over time. Second, looking toward historical patterns for definitions of progression is essentially looking at data and then designing an hypothesis to fit. Progression ought to be defined prospectively rather than retrospectively, meaning that it ought to have predictive value; if evolution is progressive, then we ought to be able to define, *a priori*, a quantity that will increase.

The definition of progression used here was purposefully chosen to deal with the above-mentioned difficulties. Thus, as it stands, it necessarily is not linked to any specific biological measurement. By the definition used here, the quantity that might increase over time need not have any obvious biological interpretation outside of the role that it plays in progressive evolution. This level of generality seems desirable if we are asking questions about the existence of such a quantity without necessarily knowing anything specific about what it might be. Such generality does mean, however, that if evolution is progressive in this sense, then the progressive trait might well be some highly complicated characteristic of the population that does not necessarily correspond to any biological attribute of an organism that is *a priori* natural. In this way, some readers might prefer to view the theorem presented here as a definition of progressive evolution rather than as a statement about the limitation of theory. In other words, we might define progressive evolution as an evolutionary process for which we could, in principle, construct a negation-complete evolutionary theory. The theorem then says that this definition is equivalent to there existing some quantity that increases over evolutionary time.

Decidability results, such as those presented here, are often prone to misinterpretation (Franzén, 2005). Therefore it is important to be clear about what the above theorem says as well as what it does not say. First, the theorem does not imply that developing a predictive theory of evolution is impossible. A very large portion of current research in evolutionary biology is directed towards developing such predictive capacity and therefore

379 the theorem takes the existence of such a theory as a starting point. The rationale is to
380 determine whether there might still be other, inherent, limits on the kinds of questions that
381 can be answered even if we are successful in pushing the development of current research in
382 this direction. The theorem demonstrates that there are such inherent limits, and in
383 essence the problem arises from a difficulty in predicting the places that evolution does not
384 go. In other words, although a predictive theory can always be used to map out the course
385 of evolution, interestingly, it cannot always be used to map out the courses that evolution
386 does not take. The theorem presented here, in effect, demonstrates that doing the latter is
387 not possible unless evolution is progressive.

388 How are these considerations to be interpreted in the context of examples like that of
389 influenza evolution discussed earlier? First, as already mentioned in that example, the
390 analysis would begin by taking what is essentially a best-case scenario, and supposing that
391 we have enough knowledge of the system to develop an open-ended model that perfectly
392 predicts (possibly in a probabilistic way) the genetic composition of the influenza
393 population in the next time step, as a function of its current composition. Then we ask, is
394 there a significant probability that another flu pandemic with the 1918 strain will ever
395 occur? The above theorem states that, even if we had such a perfect model, this kind of
396 question is unanswerable unless influenza evolution is progressive. In other words, unless
397 some characteristic of the influenza population changes directionally during evolution (e.g.,
398 some aspect of the antigenicity profile changes directionally) such a prediction will not be
399 possible. Moreover, this limitation arises because, even though we can use our perfect
400 model to map out the course of influenza evolution over time, this need not be enough to
401 map out the parts of genotype space that influenza will *not* explore.

402 The above limitations apply to predictions about the genetic evolution of the population,
403 but what if we are interested only in phenotypic predictions? For example, could we
404 predict whether or not an influenza pandemic similar in severity to that of 1918 will ever

405 occur again, regardless of which strain(s) cause the pandemic? Likewise, could we predict
406 whether or not resistance to antiviral medication will ever evolve, regardless of its genetic
407 underpinnings? If the genotype-phenotype map is one-to-one, then predicting phenotypic
408 evolution will be no different than predicting genotypic evolution. Even if many different
409 genotypes can produce the same phenotype, however, predicting phenotypic evolution still
410 involves predicting whether or not certain subsets of genotype space are visited during
411 evolution. As a result, all of the aforementioned limitations should still apply to such cases.
412 The only exception is if the genotype-phenotype map resulted in the dimension of
413 phenotype space being finite even though the dimension of the genotype space was
414 effectively infinite. Even in this case, however, the above limitations to prediction would
415 still apply unless phenotypic knowledge alone was sufficient to predict the state of the
416 population from one time step to the next (i.e., if we didn't need to consider genetic state
417 to understand evolution). While this might be possible for some phenotypes of interest, it
418 seems unlikely that it would be possible for all possible phenotypes.

419 One might argue, however, that some patterns of phenotypic evolution are very predictable.
420 For example, the application of drug pressure to populations seems inevitably to lead to
421 the evolution of resistance to the drug. How are these sorts of findings reconciled with the
422 results presented here? First, although the evolution of resistance does appear to be
423 somewhat predictable, we must distinguish between inductive versus deductive predictions.
424 One reason we feel confident about predicting the evolution of drug resistance is that we
425 have seen it occur repeatedly. Therefore, by an inductive argument we expect it to occur
426 again. Such inductive predictions are conceptually similar to extrapolating predictions
427 from a statistical model beyond the range of data available. On the other hand, deductive
428 predictions are made by deducing a prediction from an underlying set of principles or
429 mechanistic processes. In a sense, inductive predictions require no understanding of the
430 phenomenon in question whereas deductive predictions are based on some underlying
431 model of how things work. The results presented here apply solely to deductive predictions.

432 A second possibility with respect to the evolution of things like drug resistance, however, is
433 that evolution is progressive (at least at this ‘local’ scale). For example, it might well be
434 that if we formulated an accurate underlying model for how influenza evolution proceeds in
435 the presence of antiviral drug pressure, there would be some population-level quantity that
436 changes in a directional way during evolution. Indeed it seems plausible that it is precisely
437 this kind of directionality that makes us somewhat confident we can predict evolution in
438 such cases. It should be noted, however, that even if evolution is not progressive the
439 theorem presented here does not rule out the possibility that some predictions can be
440 made. For example, it is entirely possible that a theory could still be developed to make
441 negation-complete predictions about the evolution of drug resistance. The theorem simply
442 says that it will not be possible to make negation-complete predictions about any arbitrary
443 aspect of evolution unless the evolutionary process is progressive.

444 As already mentioned, all of the results presented here begin with the assumption that we
445 can develop a theory to predict evolution from one time step to the next. Whether or not
446 current theoretical approaches can be pushed the point where this is true remains a
447 separate, and open, question. There are certainly considerable obstacles to doing so unless
448 the evolutionary system of interest is very simple (e.g., Ibarra et al. (2002)). In addition to
449 the problem that historical contingencies raise, the role of uncertainty in initial conditions,
450 much like those in weather forecasting, might preclude long-term predictions (although
451 probabilistic statements might still be possible). This remains an important and active area
452 of research on which the theorem presented here offers no perspective. Rather it simply
453 reveals that, in the event that theory is eventually developed to do so, it will still face
454 inherent limitations on the kinds of questions it can answer unless evolution is progressive.

455 Although a negation-complete theory for the entire evolutionary process of interest is not
456 possible unless evolution is progressive, this also does not preclude the possibility that a
457 perfectly acceptable, negation-complete, theory might be developed for short-term and/or

458 local predictions. Indeed, just as similar inherent limitations in computability theory and
459 mathematical logic have not prevented people from making astonishing progress in these
460 areas of research, so to is the case for evolutionary biology. As mentioned in the
461 introduction, many theoretical advances have already been made by focusing on subsets of
462 the space of potential evolutionary outcomes. Continuing to push theoretical development
463 in this direction by broadening the space considered will be possible regardless of the
464 nature of the evolutionary process. The theorem does imply, however, that unless evolution
465 is progressive, it will not be possible to encompass all such developments within a single
466 unified set of principles from which all negation-complete evolutionary predictions can be
467 drawn.

468 There are some previous theoretical results in the literature that consider the extent to
469 which evolution exhibits a directional tendency and it is useful to consider how the present
470 results relate to these previous works. For example, it has been shown previously with
471 quite general stochastic models of evolution that a quantity termed ‘free fitness’ is always
472 non-decreasing during evolutionary change (Iwasa, 1988). The analysis, however, did not
473 allow for open-ended evolution because the state space was assumed to be finite, and the
474 Markov model used was (implicitly) assumed to be positively recurrent. As a result, a
475 unique stationary distribution existed and thus continual evolution was precluded.

476 It might be reasonably argued however that, although analyses such as (Iwasa, 1988) do
477 not allow for truly open-ended evolution, if the state space is large enough, and if the
478 transient dynamics are long enough, then it is effectively an open-ended model. As such,
479 should not the results with respect to free fitness still apply? In other words, does this not
480 then suggest that there is some quantity (free fitness) that increases during evolution, and
481 thus that a negation-complete theory is possible? The answer is no, and the reason is
482 subtle but important. The definition of free fitness in (Iwasa, 1988), like other quantities
483 that have been suggested to change directionally during evolution (e.g., Adami et al.

(2000)) are based on measures closely related to entropy. Importantly, the mapping between these measures of entropy and population states is not one-to-one because there are many (indeed, potentially infinitely many) biologically distinct population states that have the same value of entropy (or the same value of ‘free fitness’). As a result, even though measures like free fitness might not decrease during evolution, an indefinite amount of biologically interesting and significant evolutionary change can still occur without any change in free fitness. Roughly speaking, although measures related to things like entropy provide an interesting physical quantity that might change directionally, the relationship between entropy and quantities that are of biological interest need not be simple.

In a similar vein one might argue that, because biological evolution takes place within a physical system that is subject to the Second Law of Thermodynamics, ultimately a general measure entropy must provide a directionality to the system. Again, while this is true in terms of the system as a whole, the mapping between entropy and the population states of biological interest is not one-to-one. Thus, even though the total entropy of the entire physical system must always increase, the entropy of any component part (e.g., the biological part of interest) need not change in this way.

What do all these considerations have to say about how the process of evolution is studied, or how current theoretical research is done? Should evolutionary biologists care about such results? For instance, do the results point to new ideas that might help us do theory better? Although there is no single answer to this question, there are two points worth making in this regard. First, the distinction between open and closed-models seems like a useful, and currently somewhat under-appreciated, way to categorize models of evolution. As such it does suggest some new directions in which evolutionary theory might be taken, particularly given that open-ended models are sometimes amenable to asking novel, and potentially very important, evolutionary questions that cannot be addressed with closed models (e.g., Fontana and Buss (1994); Fontana and Schuster (1998b); Lenski et al. (1999);

510 Stadler et al. (2001); Yedid and Bell (2001); Wilke et al. (2001); Yedid and Bell (2002);
511 Lenski et al. (2003); Chow et al. (2004); Ostrowski et al. (2007)). Second, to the extent
512 that one cares about developing theory for open-ended evolutionary processes, the theorem
513 presented here then reveals that there is an inherent ‘upper bound’ on how far we can push
514 the predictive capability of such theory. In particular, although such theory opens the door
515 to asking new evolutionary questions, unless evolution is progressive, there will remain
516 some such questions that are unanswerable. Furthermore, although it will likely be difficult
517 to use the theorem as a means of proving that evolution is progressive (i.e., by developing a
518 negation-complete theory) or to use the theorem to prove that a complete evolutionary
519 theory is possible (i.e., by determining that evolution is progressive) the result does
520 nevertheless reveal that these two important, and somewhat distinct, biological ideas are
521 fundamentally one and the same thing.

522 My intention was not to imply that the theorem could be used to determine decidability
523 from knowledge of progression, or the reverse. Rather, it was to prove (within the set of
524 assumptions used) that decidability and progression can be viewed as one of the same
525 thing.

526 The theorem presented here has close ties to Gödel’s Incompleteness Theorem for
527 axiomatic theories of the natural numbers (Gödel, 1931; Nagel and Newman, 1958; Davis,
528 1965; van Heijenoort ed., 1967; Smith, 2007). An axiomatic theory consists of a set of
529 symbols, a logical apparatus (e.g., the predicate calculus), a set of axioms involving the
530 symbols, and a set of rules of deduction through which new statements involving the
531 symbols can be derived (termed ‘theorems’; Smith (2007)). Given such a system, theorems
532 can be derived through the repeated algorithmic application of the rules of deduction.

533 In the early 1900’s there was a concerted attempt to produce such an axiomatic theory
534 that was meant to represent the natural numbers, with the proviso that it yield all true
535 statements about the natural numbers, and no false ones; (Whitehead and Russell, 1910;

Smith, 2007). Gödel's Incompleteness Theorem (Gödel, 1931; Nagel and Newman, 1958; Davis, 1965; van Heijenoort ed., 1967; Smith, 2007), however, revealed that this is impossible for any axiomatic system sufficiently rich that it can make simple number-theoretic statements. For example, it shows that if the axiomatic system is rich enough that it can express the number-theoretic statement corresponding to the predicate ' $x \in R_E$ ', then it cannot produce all true number-theoretic statements and no false ones (Smith, 2007). For if it could, then it could always produce the number theoretic statement corresponding to either ' $x \in R_E$ ' or ' $x \notin R_E$ ' as a theorem, because one of the two must be true. But if it can do this, then it provides an algorithmic procedure for deciding the predicate ' $x \in R_E$ ', and we know that this is not always possible as the results presented here illustrate.

The Halting Problem from computability theory (Turing, 1936; Cutland, 1980) is also intimately related to the results presented here. As already detailed, the question of whether a population state is evolutionarily attainable is equivalent to the question of whether a given positive integer is in the range of a particular computable function. Moreover, this latter question is directly connected to the analogous question of whether a given integer is in the domain of a computable function (i.e., whether, given a particular integer input, the function returns a value in finite time). The latter problem is precisely the Halting Problem, and it is known that there is no general algorithmic procedure for solving the Halting problem for arbitrary computable functions (Turing, 1936; Cutland, 1980).

As mentioned earlier, in a very general sense, the results presented here are applicable to any system that can be faithfully described by a Markov dynamical system over an infinite set of discrete possibilities (i.e., an open-ended dynamical system). Therefore, one might ask whether there is anything in the results presented that is particular to evolution *per se*? In one sense the answer is 'no', but therein lies the power of such mathematical abstraction;

562 it reveals the underlying, key, structure of the process. Evolution will be an open-ended
563 dynamical system whenever heredity is indefinite, and it therefore shares a fundamental
564 similarity with all other processes that are also such open-ended dynamical systems.

565 At the same time, however, the results do have special significance for evolution. There are,
566 perhaps, relatively few other kinds of processes of interest that share the property of being
567 such an open-ended dynamical system in a meaningful way. For example, a great many
568 processes of interest have a relatively small space of potential outcomes, and are thus
569 clearly not open-ended. Furthermore, for those processes that are potentially open-ended,
570 it is sometimes of little theoretical interest to distinguish among all possible outcomes, and
571 therefore the space of relevant outcomes can still be relatively small. Moreover, even when
572 the space of potential outcomes of interest truly is open-ended, some processes (e.g., some
573 physical processes) obey simple enough dynamics that such negation-complete predictions
574 can readily be made (i.e., the system is ‘progressive’ in the sense considered here). Thus,
575 the limitations detailed by the theorem are of interest, primarily for those processes that
576 are both open-ended, and that are complex enough that the question of progression is
577 unresolved (Appendix 4). Evolution under indefinite heredity might be a somewhat unique
578 process in satisfying both of these criteria.

579 There are, however, other processes of interest for which such decidability results might be
580 of interest. After all, in an important sense, biological evolution is nothing more than the
581 emergent properties of physics and chemistry. In fact such limitations on theory have been
582 discussed previously, particularly as they relate to the so-called theory of everything in
583 physics (Hawking, 2002). It is probably safe to say that no general consensus on this issue
584 has yet been reached (Franzén, 2005); however, the theorem presented here has
585 implications for any physical or chemical theory that aims to explain evolutionary
586 phenomena. It demonstrates that a rational, deductive, approach to such theory will
587 necessarily face some inherent limitations on the answers that it can provide.

Acknowledgements

I thank N.H. Barton, G. Bell, S.A. Frank, S. Gandon, A. Gardner, S. Gavrillets, B. Glymour, M. Hochberg, L. Nagel, S.P. Otto, S. Proulx, A. Read, A. Rosales, P. Taylor, M. Turelli, and the Queen's Math Bio Group for inspiring conversations and feedback on the manuscript that greatly helped to clarify the results and to put them into perspective. This research was supported by an NSERC Canada Discovery Grant, an E.W.R. Steacie Fellowship, and the Canada Research Chairs Program.

References

- Adami, C., C. Ofria and T. C. Collier. 2000. Evolution of biological complexity. *P Natl Acad Sci USA* 97(9):4463–4468.
- Barton, N. H., D. E. G. Briggs, J. A. Eisen, D. B. Goldstein and N. H. Patel. 2007. *Evolution*. Cold Spring Harbor Laboratory Press.
- Bürger, R. 2000. *The Mathematical Theory of Selection, Recombination, and Mutation*. John Wiley and Sons.
- Chow, S. S., C. O. Wilke, C. Ofria, R. E. Lenski and C. Adami. 2004. Adaptive radiation from resource competition in digital organisms. *Science* 305(5680):84–86.
- Cutland, N. J. 1980. *Computability: An Introduction to Recursive Function Theory*. Cambridge University Press.
- Davis, M. 1965. *The Undecidable: Basic Papers on Undecidable Propositions, Unsolvability Problems and Computable Functions*. Raven, New York. Reprint, Dover, 2004.
- Dawkins, R. 1997. Human Chauvinism. *Evolution* 51:1015–1020.
- DeVries, H. 1904. *Species and Varieties: Their Origin by Mutation*. Chicago: Open Court.
- Ewens, W. J. 2004. *Mathematical Populations Genetics. I. Theoretical Introduction*. Springer.
- Fernando, C. and J. Rowe. 2007. Natural selection in chemical evolution. *Journal of Theoretical Biology* 247(1):152–167.
- Fisher, R. 1930. *The Genetical Theory of Natural Selection*. Clarendon Press.
- Fontana, W. and L. Buss. 1994. The arrival of the fittest - Towards a theory of biological organization. *Bulletin of Mathematical Biology* 56(1):1–64.

617 Fontana, W. and P. Schuster. 1998a. Continuity in evolution: On the nature of transitions.
618 Science 280(5368):1451–1455.

619 Fontana, W. and P. Schuster. 1998b. Shaping space: the possible and the attainable in
620 RNA genotype-phenotype mapping. Journal of Theoretical Biology 194(4):491–515.

621 Franzén, T. 2005. Gödel’s Theorem: An Incomplete Guide to Its Use and Abuse. A.K.
622 Peters Ltd.

623 Gillespie, J. H. 1984. Molecular evolution over the mutational landscape. Evolution
624 38(5):1116–1129.

625 Gödel, K. 1931. Über formal unentscheidbare Sätze der Principia Mathematica und
626 verwandter Systeme. Monatshefte für Mathematik und Physik 38:173–98.

627 Gould, S. J. 1997. Self-help for a hedgehog stuck on a molehill. Evolution 51:1020–1023.

628 Grimmett, G. and D. Stirzaker. 1992. Probability and Random Processes, second edition.
629 Oxford University Press.

630 Hawking, S. W. (2002). Gödel and the end of physics.
631 <http://www.hawking.org.uk/index.php/lectures/91>.

632 Hofbauer, J. and K. Sigmund. 1988. The Theory of Evolution and Dynamical Systems.
633 Cambridge University Press.

634 Ibarra, R., J. Edwards and B. Palsson. 2002. Escherichia coli K-12 undergoes adaptive
635 evolution to achieve in silico predicted optimal growth. Nature 420:186–189.

636 Iwasa, Y. 1988. A free fitness that always increases in evolution. Journal of Theoretical
637 Biology 135(4):265–281.

638 Joyce, P., D. R. Rokytá, C. J. Beisel and H. A. Orr. 2008. A general extreme value theory
639 model for the adaptation of DNA sequences under strong selection and weak mutation.
640 Genetics 180(4):1627–1643.

- Kimura, M. and J. Crow. 1964. The number of alleles that can be maintained in a finite population. *Genetics* 49:725–738.
- Lenski, R. E., C. Ofria, R. T. Pennock and C. Adami. 2003. The evolutionary origin of complex features. *Nature* 423(6936):139–144.
- Lenski, R. E., C. Ofria, T. C. Collier and C. Adami. 1999. Genome complexity, robustness and genetic interactions in digital organisms. *Nature* 400(6745):661–664.
- Levinton, J. 1988. *Genetics, Paleontology and Macroevolution*. Cambridge University Press.
- Lynch, M. and B. Walsh. 1998. *Genetics and Analysis of Quantitative Traits*. Sinauer and Associates.
- Manapat, M., H. Ohtsuki, R. Bürger and M. A. Nowak. 2009. Originator dynamics. *Journal of Theoretical Biology* 256(4):586–595.
- Maynard Smith, J. and E. Szathmáry. 1995. *The Major Transitions in Evolution*. Oxford University Press.
- Nagel, E. and J. R. Newman. 1958. *Gödel’s Proof*. Routledge.
- Nowak, M. A. and H. Ohtsuki. 2008. Prevolutionary dynamics and the origin of evolution. *P Natl Acad Sci USA* 105(39):14924–14927.
- Ohtsuki, H. and M. A. Nowak. 2009. Pre-life catalysts and replicators. *Proceedings of the Royal Society B: Biological Sciences* 276(1674):3783–3790.
- Orr, H. A. 1998. The population genetics of adaptation: the distribution of factors fixed during adaptive evolution. *Evolution* 52(4):935–949.
- Orr, H. A. 2002. The population genetics of adaptation: the adaptation of DNA sequences. *Evolution* 56(7):1317–1330.

664 Ostrowski, E. A., C. Ofria and R. E. Lenski. 2007. Ecological specialization and adaptive
 665 decay in digital organisms. *American Naturalist* 169(1):E1–E20.

666 Smith, P. 2007. *An Introduction to Gödel’s Theorems*. Cambridge University Press.

667 Stadler, B. M. R., P. F. Stadler, G. P. Wagner and W. Fontana. 2001. The topology of the
 668 possible: Formal spaces underlying patterns of evolutionary change. *Journal of*
 669 *Theoretical Biology* 213(2):241–274.

670 Szathmary, E. 1995. A Classification of replicators and lambda-calculus models of biological
 671 organization. *Proceedings of the Royal Society B: Biological Sciences* 260(1359):279–286.

672 Turing, A. 1936. On computable numbers, with an application to the
 673 Entscheidungsproblem. *Proceedings of the London Mathematical Society, Series 2*
 674 42:230–265.

675 van Heijenoort ed., J. 1967. *From Frege to Gödel: A Source Book on Mathematical Logic*
 676 1879-1931. Harvard University Press.

677 Wagner, G. P. and L. Altenberg. 1996. Perspective: Complex adaptations and the
 678 evolution of evolvability. *Evolution* 50(3):967–976.

679 Wagner, G. P. and P. F. Stadler. 2003. Quasi-independence, homology and the unity of
 680 type: A topological theory of characters. *Journal of Theoretical Biology* 220(4):505–527.

681 Whitehead, A. N. and B. Russell. 1910. *Principia Mathematica* v.1-3. Cambridge
 682 University Press.

683 Wilke, C. O., J. L. Wang, C. Ofria, R. E. Lenski and C. Adami. 2001. Evolution of digital
 684 organisms at high mutation rates leads to survival of the flattest. *Nature*
 685 412(6844):331–333.

686 Wright, S. 1984. *Evolution and the Genetics of Populations* v. 1-4. University of Chicago
 687 Press.

- 688 Yedid, G. and G. Bell. 2001. Microevolution in an electronic microcosm. *American*
689 *Naturalist* 157(5):465–487.
- 690 Yedid, G. and G. Bell. 2002. Macroevolution simulated with autonomously replicating
691 computer programs. *Nature* 420(6917):810–812.
- 692 Yedid, G., C. A. Ofria and R. E. Lenski. 2008. Historical and contingent factors affect
693 re-evolution of a complex feature lost during mass extinction in communities of digital
694 organisms. *J Evolutionary Biol* 21(5):1335–1357.
- 695 Yedid, G., C. A. Ofria and R. E. Lenski. 2009. Selective Press Extinctions, but Not
696 Random Pulse Extinctions, Cause Delayed Ecological Recovery in Communities of
697 Digital Organisms. *American Naturalist* 173(4):E139–E154.

698 Figure 1: A schematic representation of the coding of population states, and the theorem.
 699 Middle irregular shape represents the space of population states, S , with four states
 700 depicted (the ovals). Roman numerals indicate the time when each state is visited during
 701 evolution (silver-shaded state, $s = \{T, T, T\}$, is never visited). Vertical ovals on right and
 702 left represent two different codings by the positive integers, along with their respective
 703 evolutionary mappings, $\phi_E(n)$ and $\phi_{\hat{E}}(n)$, over the first three time steps. If evolution is
 704 progressive, then Coding 2 is possible, and the theorem then says we can ‘decide’ any
 705 population state, $s \in S$. For example, we can decide state ‘T,T,T’ by finding its code (i.e.,
 706 ‘1’), and then iterating the map, $\phi_{\hat{E}}(n)$, until we obtain an output greater than ‘1’ (this
 707 occurs at time step 1 because $\phi_{\hat{E}}(1) = 2$). If ‘1’ has not yet been visited by this time, it
 708 never will be. Conversely, if all population states are decidable, then under Coding 1 we
 709 can apply the algorithm provided in Part 2 of the theorem’s proof to obtain Coding 2,
 710 thereby demonstrating that evolution is progressive.

Appendices

1 Theory

The term ‘theory’ is used in a technical sense. A theory consists of a set of symbols that constitute the language of the theory, a set of premises which are taken as given, and a set of rules of inference (Smith, 2007). The symbols represent certain components of reality, and the premises constitute statements about reality through the interpretation of the symbols. The rules of inference then constitute valid ways of deducing new statements about the symbols of the language, and thus through interpretation, new statements about reality. Thus, within such a theory, statements are derived by taking some premise(s), and applying the rules of inference.

Statements derived through a series of deductive arguments using the rules of inference are referred to as theorems of the theory. The result of the main text is valid for any evolutionary theory whose theorems are recursively enumerable (Appendix 2); i.e., any theory whose theorems can be derived through the use of a finite (but possibly large) number of mechanical, or algorithmic, steps (e.g., as laid out in the rules of inference; Appendix 2). This is clearly true for any such theory based on computation, since computers do nothing more than mechanically follow rules (Cutland, 1980). It is also true for any axiomatic theory, since the theorems of any such theory can be derived simply by applying the mechanical rules of inference to the axioms (Smith, 2007).

A great deal of current quantitative theory in evolutionary biology fits the above template. For example, current theory often abstracts reality mathematically by assigning formal symbols to things like allele frequencies and population sizes. A set of premises is then taken, for example, by formalizing an hypothesis about how genotypic fitnesses are determined. Next, a finite number of applications of ‘rules of inference’ are used (e.g., the

application of certain mathematical operations) in order to derive statements about the formal symbols of this theory. Finally, these symbolic statements are then interpreted again in terms of their biological meaning, and hence predictions about evolution are made (Fig. S1).

Figure S1: A schematic representation of the relationship between the biological process of evolution and theory. The example given illustrates classical population-genetic theory. A formal system is created to represent elements of evolution (e.g., $p(t)$ represents the number of the blue genotype at time t). A set of premises is specified (e.g., initial genotype numbers, how genotypic fitnesses are determined, etc. - this is embodied by the mapping F). Rules of deduction are then followed (e.g., repeated application of the mapping F) to obtain new statements about elements of the formal theory (e.g., $p(1), p(2), p(3)$ etc.). These new elements are then interpreted in terms of evolution (e.g., as predictions about genotype numbers at future times).

2 Some results from computability theory

A function is computable if it can be evaluated by an Unlimited Register Machine (URM) in a finite numbers of steps (Cutland, 1980). The Church-Turing Thesis states that any function we might view as being evaluated through a mechanical procedure can be evaluated by a URM (Cutland, 1980). Thus, given the Church-Turing Thesis, the easiest way to ascertain whether something is computable is to consider whether a computer could be programed to do it in such a way that an output is guaranteed, in a finite (but possibly very large) number of steps.

Definition: A function is total if it is computable over all natural numbers.

Definition: A function is partial if it is computable only over some (nonempty) subset of

758 the natural numbers.

759 **Definition:** A set is denumerable if there exists a bijection between it and the natural
760 numbers.

761 **Definition:** A set is effectively denumerable if this bijection, and its inverse, are
762 computable.

763 **Definition:** The characteristic function of a set of natural numbers, A , is

$$c_A(n) = \begin{cases} 1 & \text{if } n \in A \\ 0 & \text{if } n \notin A \end{cases} \quad (1)$$

764 **Definition:** The predicate ' $n \in A$ ' is decidable if its characteristic function is computable.

765 **Definition:** The set A is recursive if the predicate ' $n \in A$ ' is decidable.

766 **Definition:** The partial characteristic function of a set of natural numbers, A , is

$$\bar{c}_A(n) = \begin{cases} 1 & \text{if } n \in A \\ \text{undefined} & \text{if } n \notin A \end{cases} \quad (2)$$

767 **Definition:** The predicate ' $n \in A$ ' is partially decidable if its partial characteristic
768 function is computable for $n \in A$.

769 **Definition:** The set A is recursively enumerable (denoted r.e.) if the predicate ' $n \in A$ ' is
770 partially decidable.

771 Note that every recursive set is r.e. but not vice versa. Furthermore, a set A is recursive if,
772 and only if, both A and its complement A^c are r.e.. Finally, note that any finite set of
773 numbers is recursive (Cutland, 1980).

774 The following concepts and notation will also prove useful:

775 First, because any computable function can be evaluated through a series of steps, we can
776 define $c_A^o(n)$ as the value of $c_A(n)$ after the o^{th} step in its evaluation. In particular, $c_A^o(n)$
777 evaluates to ‘null’ if it has not returned a value by the o^{th} step.

778 Second, a standard result from computability theory demonstrates that there exists a
779 computable bijection between $\mathbb{N}^+$ and $\mathbb{N}^+ \times \mathbb{N}^+$ (Cutland, 1980). We will denote this
780 mapping by $B : n \mapsto (T_1(n), T_2(n))$.

781 Third, the notion of an ‘unbounded search’ is central in computability theory. In
782 particular, it is standard to use the notation $\mu y(f(y) = k)$ to denote ‘the smallest value of
783 y such that $f(y) = k$ ’.

784 Fourth, a fundamental theorem of computability theory demonstrates that the set of all
785 computable functions is denumerable (Cutland, 1980). Thus, we can use $\phi_k(n)$ to denote
786 the k^{th} computable function, and R_k and D_k as its range and domain respectively. We will
787 also make use of the notation $R_k(n) = \{x : \phi_k(i) = x, i \leq n\}$. In other words, if $\phi_k(n)$ is
788 evaluated for increasing values of n , then $R_k(n)$ is the subset of the range of $\phi_k(n)$ that has
789 been visited by step n . This is clearly computable for any n if $\phi_k(n)$ is total.

790 Finally, notice that it was implicitly assumed that the mapping, G corresponding to the
791 evolutionary process is computable, and thus $E(n)$ is a computable function. Thus, the
792 evolutionary process is, in an important way, nothing other than computation. Although it
793 is not practically feasible to verify or refute this assumption for most evolutionary systems,
794 there are very good reasons to expect that this assumption is reasonable. First, if we are

willing to view the processes occurring in our biological system as being purely ‘mechanical’, then we can appeal to the Church-Turing Thesis to argue that G must thereby be computable. Second, the use of the term ‘evolution’, as a process, should not be restricted to a particular instantiation of this process, as for example occurs in carbon-based life. For example, there are very good reasons to think that the processes occurring in *in silico* evolution are fundamentally the same as those occurring in biological evolution. As such these would clearly be computable. Finally, even if biological evolution isn’t formally computable (i.e., it is not mechanical) we nevertheless usually proceed by assuming that it can be modeled using computation.

3 The set of population states is effectively denumerable

Here we prove that the set of possible population states is effectively denumerable; i.e., that there exists a computable bijection between the population states and the positive integers with a computable inverse. Such sets are also called effectively denumerable.

Proof: We simply need to demonstrate an effective procedure (i.e., a computable procedure) for both encoding and decoding the population states into positive integers. Let M be the maximum possible population size (a positive integer). Each of the M ‘slots’ is either vacant, or filled by an individual that is completely characterized by its DNA sequence. Furthermore, we can set A=0, C=1, G=2, T=3, and then read the DNA sequence from its 5’ to 3’ end, thereby establishing a unique characterization of each slot in the population.

(A) Encoding: For each of the M slots calculate a numeric code as follows: Reading the DNA from its 5’ to 3’ end, for the n^{th} base, take the n^{th} prime number and raise it to the power corresponding to this base as listed above. Multiply all these numbers together.

This gives a unique number for each distinct DNA sequence, and thus the mapping is injective. Furthermore, since all positive integers greater than or equal to 2 have a unique prime factorization, all such integers correspond to a DNA sequence. Thus, if we code the state ‘vacant’ with the number 1, the mapping is surjective as well. Furthermore, this procedure is computable for any piece of DNA. This shows that there is a computable encoding for each slot, and since the population is simply the union of a finite number of such slots, the population state has a computable encoding as well. In particular, the coding of each slot locates a point in $\mathbb{N}^+ \times \cdots \times \mathbb{N}^+$ (where $\mathbb{N}^+$ appears M times) that can be uniquely identified by its indices. One can then cycle through all possible indices as follows: start with all indices that sum to 1, then those that sum to 2 etc. This is computable, and for each instance we simply assign a code number in increasing order.

(B) Decoding: For any given code number, cycle through the sets of indices as above, stopping once the code number is reached, and determine those indices. Once these indices have been obtained, one can determine their corresponding DNA through their prime factorization.

4 Some additional technical information about the theorem

The theorem of the text would be of little interest if it were never possible for ‘ $x \in R_E$ ’ to be undecidable. It is well-known in computability theory that there exist computable functions for which such predicates are undecidable ((Cutland, 1980); Appendix 4), but the evolutionary process considered represents a special kind of computable function. In particular, it must satisfy the mapping $\phi_k(n+1) = G(\phi_k(n))$ for all n , where $G()$ is a computable function with appropriate domain. The subset of computable functions satisfying this relation will be referred to as Markov, total, computable functions.

842 This section presents a series of three lemmas that, together, demonstrate that there do in
843 fact exist Markov computable functions for which ‘ $x \in R_E$ ’ is undecidable (see also
844 Cutland (1980); Smith (2007)). In such cases, the set of evolutionarily attainable states,
845 R_E will be called ‘recursively enumerable’ (r.e.; because ‘ $x \in R_E$ ’ is always at least
846 partially decidable for Markov computable functions). On the other hand, if ‘ $x \in R_E$ ’ is
847 decidable, then R_E is said to be ‘recursive’ (Appendix 2 and Appendix 4).

848 *Lemma 1: A set of numbers is recursively enumerable if, and only if, it is the range of*
849 *some total, computable, function. Note: we could relax the ‘total’ requirement without much*
850 *change.*

851 Proof: (i) A r.e. $\Rightarrow$ ‘ A is the range of a total computable function’

852 Given A is r.e., the partial characteristic function of A is computable; i.e.,

$$\bar{c}_A(n) = \begin{cases} 1 & \text{if } n \in A \\ \text{undefined} & \text{if } n \notin A \end{cases} \quad (3)$$

853 is computable. Now first choose an $a \in A$. This is a computable operation since we can
854 simply use the bijection $B : n \mapsto (T_1(n), T_2(n))$ to evaluate $\bar{c}_A^{T_2(n)}(T_1(n))$ for increasing n
855 until it returns a value of 1, and then identify the corresponding value $T_1(n)$. Next, we can
856 define the computable function

$$g(x, o) = \begin{cases} x & \text{if } \bar{c}_A^o(x) = 1 \\ a & \text{otherwise} \end{cases} \quad (4)$$

857 Then, again we can use the computable bijection $B : n \mapsto (T_1(n), T_2(n))$ to define
858 $f(n) = g(T_1(n), T_2(n))$. This is a total computable function with range equal to A .

859 (ii) ' R_k is the range of a total computable function' $\Rightarrow R_k$ r.e.

860 Consider the total function $\phi_k(n)$. We can then construct the computable partial
861 characteristic function for R_k as follows: For any input value, x , output the value 1 after
862 evaluating $\mu i(\phi_k(i) = x)$.

863 Q.E.D.

864 Given Lemma 1, we can then prove the following, second, lemma;

865 *Lemma 2: There exists total computable functions whose ranges are r.e. but not recursive.*

866 Using Lemma 1, we can prove Lemma 2 by proving that there exist sets that are r.e. but
867 whose complements are not r.e.

868 *Proof Sketch (by construction) see Smith (2007):*

869 We will demonstrate that $K = \{n : n \in R_n\}$ is one such set. It is clear, therefore, that
870 other such sets can be constructed as well.

871 First it can be proven that K^c is not r.e. using Cantor's diagonal argument (e.g., see Smith
872 (2007)). In particular, since all r.e. sets are the range of some computable function, and
873 since the computable functions are denumerable, the set of all r.e. sets is denumerable. So
874 we simply need to construct a set that is not in this list. Choosing numbers n such that
875 $n \notin R_n$ satisfies this property, and this is exactly K^c .

876 All that remains then is to show that K is r.e. As with characteristic functions, all
877 computable functions are evaluated through a series of operations for each input, and
878 therefore we can consider the σ^{th} operation of any computable function. Therefore, define

$$g(x, o, n) = \begin{cases} \phi_n(x) & \text{if } \phi_n(x) \text{ halted by operation } o \text{ in its evaluation} \\ n + 1 & \text{otherwise} \end{cases} \quad (5)$$

879 This is a computable function. Now we can use the bijection $B : n \mapsto (T_1(n), T_2(n))$ to
880 define $f(z, n) = g(T_1(z), T_2(z), n)$. This is also computable, and for any given n and z it
881 outputs either $n + 1$ or else an element of R_n . We can then construct the computable
882 partial characteristic function for K as follows: For any input value, n , output the value 1
883 after evaluating $\mu z(f(z, n) = n)$.

884 These results show that there exist computable functions whose ranges are r.e. but not
885 recursive. Note that some such functions might have the same output values for more than
886 one value in their domain, but these cannot be Markov computable functions. The reason
887 is simply that the mapping G ensures that, if R_E is infinite, then $\phi_E(n)$ can never repeat
888 itself as n increases (see Lemma 1, Appendix 5). Therefore, we still need to demonstrate
889 that, even if we restrict attention to Markov computable functions, some such functions
890 have r.e. ranges that are not recursive. This is done in the third lemma:

891 *Lemma 3: For every total computable function having a range that is r.e. but not recursive,*
892 *there exists a total computable Markov function with the same range.*

893 Proof: Suppose that $\phi_k(n)$ is total and has an r.e. range that is not recursive (and thus R_k
894 is infinite). Define the computable function $\phi_{\hat{k}}(n) = \phi_k(z(n))$, where
895 $z(n) = \mu i(\phi_k(i) \notin R_k(n - 1))$. It is clear that $\phi_{\hat{k}}(n)$ is a total, computable function with
896 range R_k . Now we simply need to show that $\phi_{\hat{k}}(n + 1) = G(\phi_{\hat{k}}(n))$ for all n for some
897 computable $G()$. By construction we can see that the computable function
898 $G(y) = \phi_{\hat{k}}(\mu z(\phi_{\hat{k}}(z) = y) + 1)$ works, where its domain is R_k . This function takes a state y ,
899 finds the unique time at which this state occurs (i.e., $\mu z(\phi_{\hat{k}}(z) = y)$ - this is computable),

900 and then adds 1. The resulting value is then used in the function $\phi_{\hat{k}}(n)$ to compute the
901 state in the next time step. In particular, we can see that $G(\phi_{\hat{k}}(n)) = \phi_{\hat{k}}(n+1)$.

902 Q.E.D.

903 5 Continuous Time & Stochasticity

904 For simplicity of exposition, all results of the main text have assumed that the evolutionary
905 process is deterministic and that generations are discrete. Here we show that an analogous
906 theorem holds if we relax these restrictions.

907 To begin, it is easy to see that the assumption of discrete generations is immaterial. In
908 particular, if we take generations to be continuous, then we can suppose that, at any
909 instant in time, only a single event is possible (e.g., individual birth or death). Thus,
910 because the state space is discrete, we can simply view the continuous-time process as one
911 in which discrete events occur at points in time that need not be uniformly spaced.

912 Allowing for stochasticity requires more work. If the evolutionary process is deterministic,
913 then there is a single population state possible for each point in time, n . In the analysis of
914 this case, we supposed that we had complete knowledge, not only of the evolutionary
915 mapping, G and its initial condition, but of the solution to this mapping, $\phi_E(n)$ as well (and
916 it is a total, computable, function).

917 Now there will be uncertainty in what the population state will be at time n , and in fact
918 there will potentially be several different states that the population might attain at n .
919 Some of these might be more likely than others in that, if we replayed the evolutionary
920 process multiple times, certain states might arise more often than others. Thus we might
921 imagine a probability distribution over the set of positive integers at each time step, n . By

922 analogy with the deterministic case, we make a Markov assumption, meaning that the
923 probability distribution on the population states at any given time, n , depends only on the
924 population state in the previous time, $n - 1$. In other words, there is some mapping, H ,
925 from current population state to the probability distribution over the population states in
926 the next time period. The solution of this mapping (given an initial condition) then gives
927 the probability distribution over the states at each point in time.

928 Just as with the deterministic case, we suppose that we have complete knowledge of the
929 solution of this evolutionary process in the following sense: at any time n , we have a total,
930 computable function that tells us simply the set of states, at that time, that have positive
931 support. Thus, we have a total, computable, set-valued function $\tilde{\phi}_E(n)$ that gives the set of
932 “feasible” states at time n . The ‘tilde’ signals that this function is now a set-valued
933 function, rather than an integer-valued one. And again the goal of a negation-complete
934 theory would then be to decide whether any given state lies within the set of feasible states
935 or not.

936 One objection to this formulation is that we might expect all states have some nonzero
937 probability, even if it is vanishingly small. As such, under this definition all states would
938 then be trivially feasible. There are at least two potential responses to this objection.
939 First, while it is true that many models of evolution assume that all states have nonzero
940 probability (e.g., many stochastic models of mutation-selection balance, including those
941 with an infinite number of different alleles; Kimura and Crow (1964)), this is usually
942 because they are ‘closed’ models in the sense described earlier. In particular they often
943 assume, for mathematical convenience, that the stochastic process is irreducible and
944 positively recurrent. This then implies that a unique stationary distribution exists
945 (Grimmett and Stirzaker, 1992) and thereby rules out the possibility of open-ended
946 evolution. Although it is possible to develop a model for open-ended evolution that still
947 has nonzero probability for all states, it is not obvious that this need be true of real

open-ended evolution. For example, out of the effectively infinite number of different nucleotide combinations that could make up a genotype, we might expect at least some of these to be truly lethal. On a more practical level, given the analysis presented here it seems reasonable to expect that a similar theorem could be proved if we instead defined a state as being feasible if it occurred with some probability greater than a small threshold value, $\epsilon > 0$. At this point, however, such a theorem remains conjecture.

Given that all of our considerations with respect to computability have been restricted to integer-valued functions, we now need to make the notion of computability of $\tilde{\phi}_E(n)$ more precise. The set-valued function $\tilde{\phi}_E(n)$ can be thought of as consisting of two separate computable functions, each of which is an integer-valued function and so fits within the notions of computability already discussed. The first function is simply a computable function $\phi_E(i)$ as before, whose range is now thought of as the set of feasible population states. The argument i here is now no longer meant to be evolutionary time, however, but rather is simply an index whose meaning is described below. The second computable function we denote by $\phi_{E^*}(n)$, and it specifies the number of feasible population states in generation n in the following way: the set of all feasible population states at time 1; i.e., $\tilde{\phi}_E(1)$ is given by $\{\phi_E(1), \phi_E(2), \dots, \phi_E(k_1)\}$, where $\phi_{E^*}(1) = k_1$. Likewise, $\tilde{\phi}_E(2) = \{\phi_E(k_1 + 1), \dots, \phi_E(k_1 + k_2)\}$, where $\phi_{E^*}(2) = k_2$, and so on. In this way, we can apply the same notions of computability to the set-valued function $\tilde{\phi}_E(n)$ by applying them to its component, integer-valued, functions $\phi_E(i)$ and $\phi_{E^*}(n)$. We will assume that the set $\tilde{\phi}_E(n)$ is finite for all n , which guarantees that it be computable. Nevertheless, it seems reasonable to expect that some formulations in which this set is infinite would still be computable, and thus would still fit within the results that follow.

As in the deterministic case, we must also specify the initial conditions, in addition to the mapping, H . Then, in terms of the mapping, H , if $x \in \tilde{\phi}_E(n)$ is a feasible population state at time n , the set of feasible population states at time $n + 1$ is given by

974 $\tilde{\phi}_E(n+1) = \bigcup_{x \in \tilde{\phi}_E(n)} \text{support} H(x)$, where $\text{support} H(x)$ denotes the set of states for which
975 $H(x)$ has positive support. The range of $\tilde{\phi}_E(n)$ is the set of all states that are feasible at
976 some time (i.e., it is the range of $\phi_E(i)$). Likewise, a state is evolutionarily attainable if
977 there is some time for which it is feasible. A complete evolutionary theory is one for which
978 the predicate ' $x \in R_E$ ' is decidable; i.e., if, given any population state, we can decide
979 whether it is feasible at some time.

980 The same definition of progressive evolution can be used in both the deterministic and
981 stochastic cases. To specify this precisely, we need the following Lemmas;

982 *Lemma 1: In the deterministic case, a new state is visited every time step if, and only if,*
983 *evolution is unbounded (i.e., R_E is infinite)*

984 *Lemma 2: In the stochastic case, at least one new state is feasible every time step if, and*
985 *only if, evolution is unbounded (i.e., R_E is infinite)*

986 Proof is given of Lemma 2 only (Lemma 1 can be proven in an analogous fashion). We
987 note that, in the remainder of this section, we use the notation $R_E(n)$ to denote the set of
988 population states that have been visited (i.e., feasible) by step n of the set-valued function,
989 $\tilde{\phi}_E(n)$ (i.e., not step n of $\phi_E(n)$). Equivalently, it denotes the range of $\phi_E(i)$ visited by step
990 $i = k_1 + k_2 + \dots + k_n$.

991 Proof:

992 'At least one new state is feasible each time step' $\Rightarrow$ 'Evolution unbounded'

993 This direction of the implication is obvious since, if at least one new state is feasible each
994 time step, then the fact that $\tilde{\phi}_E(n)$ is total implies that R_E is infinite.

995 'Evolution unbounded' $\Rightarrow$ 'At least one new state is feasible each time step'

996 Contrary to the assertion, suppose instead that R_E is infinite but that there is some time,
 997 n^* at which no new state is feasible. In other words, for some time n^* , the set $\tilde{\phi}_E(n^*)$
 998 satisfies $\tilde{\phi}_E(n^*) \subseteq R_E(n^* - 1)$. The set of feasible states in the next time step is then given
 999 by $\tilde{\phi}_E(n^* + 1) = \bigcup_{x \in \tilde{\phi}_E(n^*)} \text{support} H(x)$. Furthermore, for each element, $x \in \tilde{\phi}_E(n^*)$,
 1000 $\exists n_x < n^*$ such that $x \in \tilde{\phi}_E(n_x)$ (from the hypothesis that $\tilde{\phi}_E(n^*) \subseteq R_E(n^* - 1)$).
 1001 Therefore, for each element, $x \in \tilde{\phi}_E(n^*)$, we have that $\text{support} H(x) \subseteq \tilde{\phi}_E(n_x + 1)$, where
 1002 $n_x < n^*$. Thus, we have

$$\tilde{\phi}_E(n^* + 1) = \bigcup_{x \in \tilde{\phi}_E(n^*)} \text{support} H(x) \quad (6)$$

$$\subseteq \bigcup_{x \in \tilde{\phi}_E(n^*)} \tilde{\phi}_E(n_x + 1) \quad (7)$$

$$\subseteq R_E(n^* - 1). \quad (8)$$

1003 Hence, by induction, $R_E \equiv R_E(n^* - 1)$, which is finite, yielding a contradiction.

1004 Q.E.D.

1005 Notice that, in the deterministic case, when evolution is unbounded the computable
 1006 function $\phi_E(i)$ never repeats a previously attained value as i increases (Lemma 1 above).
 1007 In the stochastic case, however, even when evolution is unbounded, $\phi_E(i)$ *can* repeat
 1008 previously attained values as i increases. The key connection between the two cases is that,
 1009 in the stochastic case, $\phi_{E^*}(n)$ is such that, when the outputs of $\phi_E(i)$ are grouped into
 1010 their corresponding evolutionary generations, each such grouping always contains at least 1
 1011 new feasible state (Lemma 2 above).

1012 Now, returning to the proof of the theorem, in the deterministic case, Lemma 1 shows that
 1013 a new population state is visited at every time step. And if evolution is progressive, then

1014 there is some way to recode the populations states such that, the code number of these new
1015 states that are visited over time increases. Likewise, Lemma 2 shows that at least one new
1016 population state becomes feasible at every time step, although some visited population
1017 states might have been visited previously as well. Nevertheless, we still say that evolution
1018 is progressive if there is some way to recode the populations states such that, the code
1019 number(s) of the new states that become feasible each time step, increases with time.
1020 Formally, if we define $\sigma_{\hat{E}}(n) = R_{\hat{E}}(n) \setminus R_{\hat{E}}(n-1)$ as the set of newly feasible states in
1021 generation n , and $\min \sigma_{\hat{E}}(n)$ as the smallest of these, then evolution is progressive if there
1022 exists a computable bijection, $\hat{C}$, between the positive integers and the population states,
1023 such that $\min \sigma_{\hat{E}}(n+1) > \min \sigma_{\hat{E}}(n)$ for all n . Since the set $R_{\hat{E}}(n)$ is finite and
1024 computable for all n , $\min \sigma_{\hat{E}}(n)$ is a total computable function.

1025 The proof of the theorem then goes through as follows:

1026 *Theorem: $x \in R_E$ is decidable (i.e., R_E is recursive) if, and only if, there exists a*
1027 *computable one-to-one coding of the population states by positive integers, $\hat{C}$, such that, for*
1028 *the corresponding description of the evolutionary process, $\tilde{\phi}_{\hat{E}}(n)$,*
1029 *$\min \sigma_{\hat{E}}(n+1) > \min \sigma_{\hat{E}}(n)$ for all n .*

1030 Proof:

1031 Part 1: $\exists \hat{C}$ s.t. $\min \sigma_{\hat{E}}(n+1) > \min \sigma_{\hat{E}}(n) \forall n \Rightarrow R_E$ recursive

1032 By hypothesis there exists a computable bijection $\hat{C}$ such that $\min \sigma_{\hat{E}}(n+1) > \min \sigma_{\hat{E}}(n)$
1033 for all n . Now for any population state, x , in the original coding, let $\hat{x}$ be the corresponding
1034 code under bijection $\hat{C}$. Define $z(\hat{x}) = \mu i(\min \sigma_{\hat{E}}(i) \geq \hat{x})$. Clearly ' $\hat{x} \in R_{\hat{E}}(z(\hat{x}))$ ' is
1035 decidable since $R_{\hat{E}}(z(\hat{x}))$ is finite and enumerable. Furthermore $\hat{x} \in R_{\hat{E}}(z(\hat{x})) \Leftrightarrow \hat{x} \in R_{\hat{E}}$
1036 owing to the progressive nature of evolution. Therefore, ' $\hat{x} \in R_{\hat{E}}$ ' is decidable as well.
1037 Finally, using S denote the set of population states that are evolutionarily attainable, we
1038 have that $\hat{x} \in R_{\hat{E}} \Leftrightarrow \hat{C}^{-1}\hat{x} \in S \Leftrightarrow C\hat{C}^{-1}\hat{x} \in R_E$. Noting that, by definition, $x = C\hat{C}^{-1}\hat{x}$,

1039 we obtain $\hat{x} \in R_{\hat{E}} \Leftrightarrow x \in R_E$. Thus, ' $x \in R_E$ ' is decidable as well.

1040 Part 2: R_E recursive $\Rightarrow \exists \hat{C}$ s.t. $\min \sigma_{\hat{E}}(n+1) > \min \sigma_{\hat{E}}(n) \forall n$

1041 We can construct the required computable bijection to show that evolution is progressive
1042 as follows.

1043 Since R_E is recursive, we know that ' $x \in R_E$ ' is decidable. So take the population states, x ,
1044 in order and go down the list using the following algorithm:

1045 (i) if $x \notin R_E$ and it is the k^{th} such state up to that point, return the k^{th} odd number.

1046 (ii) if $x \in R_E$, and if it has not yet been assigned a new code number, do the following:

- 1047 • calculate $\mu i(x \in \tilde{\phi}_E(i))$ (i.e., the first time that x becomes feasible).
- 1048 • calculate $\sigma_E(i)$, the entire set of newly feasible states at i .
- 1049 • using the notation $|A|$ to denote the cardinality of A , assign codes to all of the $|\sigma_E(i)|$
1050 elements in $\sigma_E(i)$, by starting with the $|R_E(i-1)| + 1$ even number, up to the $|R_E(i)|$
1051 even number, in any order.
- 1052 • move on to the next state in the list.

1053 Thus, $R_{\hat{E}}$ is again the set of even numbers, and the new states that are feasible each time
1054 step always have larger code values as time increases. In particular, using $\hat{C}C^{-1}$ to denote
1055 the algorithm described above in points (i) and (ii), where C^{-1} is the inverse mapping of
1056 the coding that generated x (i.e., it takes code x and returns the corresponding population
1057 state, s), we have $\min \sigma_{\hat{E}}(n+1) = \min \hat{C}C^{-1}\sigma_E(n+1) = 2|R_E(n) + 1|$. The last equality
1058 follows from the fact that $\hat{C}C^{-1}\sigma_E(n+1)$ determines the first time that each element of
1059 $\sigma_E(n+1)$ occurs (which is $n+1$ for all such elements by definition), and then assigns the
1060 codes $2|R_E(n) + 1|$ up to $2|R_E(n+1)|$ for these elements. The minimum of these codes is,

of course, $2|R_E(n) + 1|$ giving $\min \sigma_{\hat{E}}(n + 1) = 2|R_E(n) + 1|$. As a result,
 $\min \sigma_{\hat{E}}(n + 1) > \min \sigma_{\hat{E}}(n)$ because $|R_E(n)|$ is strictly increasing with n (from Lemma 2).

Q.E.D.

6 Effectively Infinite Systems

The simplified system of evolution considered in the main text assumes that the space of potential population states is infinite, and focuses on unbounded evolution (i.e., $|R_E| = \infty$). One might argue, however, that any real system of evolution is necessarily finite, if only because of a potential limit to the constituent elements of the genetic material. There are two potential responses to this objection. First, on a philosophical level, although any particular evolutionary system might be finite, one might nevertheless want evolutionary theory to stand abstractly, independent of any particular instantiation of an evolutionary dynamic. This is very much analogous to the fact that, in the context of number theory, although one necessarily only ever has to deal with a finite number of things that require counting, we nevertheless desire an abstract theory of numbers that does not presuppose any finite limitations. And just as such a negation-complete theory of numbers is not possible (Gödel, 1931; Nagel and Newman, 1958; Davis, 1965; van Heijenoort ed., 1967; Smith, 2007), neither is one for evolutionary biology unless evolution is progressive.

Second, on a more practical level, it is clear that the digital nature of heredity offered by DNA/RNA makes such systems effectively infinite in that the number of possible population states is enormous. The remainder of this section makes the notation of effectively infinite precise. For simplicity, the focus below is on the deterministic system.

Recall that, in the $|R_E| = \infty$ case, a function is computable (and total) if it can be evaluated in a finite number of steps, for any input (Cutland, 1980) (Appendix 2). Thus

1084 the predicate ' $x \in R_E$ ' is decidable if its characteristic function can be evaluated, for any
 1085 input value x , in a finite number of steps. Likewise, the mapping $\hat{C}$ of the theorem is
 1086 computable if, for any input, it returns a code number in a finite number of steps.

1087 When $|R_E| < \infty$, however, the predicate ' $x \in R_E$ ' is always decidable because we can
 1088 always carry out a complete cataloguing of R_E in a finite number of steps. We simply need
 1089 to successively evaluate $\phi_E(n)$ for increasing values of n . According to Lemma 1 of
 1090 Appendix 5, because R_E is finite, we will eventually obtain a value that has previously
 1091 been visited, and from that point onward the system will then simply revisit previously
 1092 visited states.

1093 Although these observations are formally correct, they nevertheless fail to capture the
 1094 important consequences of digital inheritance in finite systems. In particular, the natural
 1095 analogue of computability for such finite systems in the context of indefinite heredity is not
 1096 the requirement that an output be obtained in a finite number of steps. Rather, it is that
 1097 an output be obtained in a finite number of steps, *and that this number of steps not exceed*
 1098 *some finite bound that is independent of the size of the state space, $|R_E|$* . For example, with
 1099 this definition for finite state spaces, the predicate ' $x \in R_E$ ' would be decidable if its
 1100 characteristic function can be evaluated in a finite number of steps, and if this number
 1101 never exceeds some finite bound that is independent of $|R_E|$. Thus, regardless of the size of
 1102 $|R_E|$, we are guaranteed to never need more than a fixed number of computational steps.

1103 To formalize these ideas, we need to be precise about what it means to consider state
 1104 spaces of different sizes, $|R_E|$. We do this as follows. First, consider the infinite state space
 1105 situation used in the main text, where $\phi_E(n)$ denotes the computable function
 1106 corresponding to the evolutionary process. Next, define the finite state space process by a
 1107 computable function, $F_E^\eta(n)$, where $n = \eta + 1$ is the first time at which a previously visited
 1108 population state is re-visited, and where $F_E^\eta(n) = \phi_E(n)$ for all $n \leq \eta$. Note that we have
 1109 $\eta = |R_E|$, and thus η is the state space size. In this way, any given finite state space

1110 process is identical to the reference infinite state space process, $\phi_E(n)$, over time until the
 1111 point $\eta + 1$ at which the finite process begins to revisit previously visited states. Thus we
 1112 can consider state spaces of different sizes, η , with the limiting case of $\eta \rightarrow \infty$
 1113 corresponding to the infinite state space of the main text. We have the following revised
 1114 definitions for the finite case:

1115 **Definition:** The predicate ' $x \in R_E$ ' is *decidable if, for any input x , there exists a $T < \infty$
 1116 such that the characteristic function $c_{R_E}(x)$ can be evaluated in no more than T steps,
 1117 where T is independent of η (i.e., independent of system size).

1118 **Definition:** A one-to-one mapping of the population states by the positive integers, $\hat{C}$, is
 1119 *computable if, for any input there exists a $T < \infty$ such that the mapping can be
 1120 evaluated in no more than T steps, where T is independent of η .

1121 The main theorem of the text can again be seen to hold when $|R_E| < \infty$ if we use the
 1122 above definitions. In particular,

1123 *Theorem: ' $x \in R_E$ ' is *decidable if, and only if, there exists an *computable one-to-one*
 1124 *coding of the population states by a subset of the positive integers, $\hat{C}$, such that the*
 1125 *corresponding description of the evolutionary process, $F_E^\eta(n)$, satisfies $F_E^\eta(n+1) > F_E^\eta(n)$*
 1126 *for all $n \leq \eta$.*

1127 Notice that there is one difference from the main theorem of the text; namely, the altered
 1128 characterization of progressive evolution. Now, because R_E is finite, we say that evolution
 1129 is progressive if there is some quantity that increases over time before the process begins to
 1130 repeat. Also note that, in addition to the altered definition of 'computable' and 'decidable'
 1131 in the statement of the theorem, all other instances of computability use this altered
 1132 definition as well.

1133 Only a sketch of a formal proof is given for this modified theorem because it is similar that

1134 of the main text. Recall that $F_E^\eta(n)$ denotes the computable function corresponding to the
 1135 finite evolutionary system of interest.

1136 Proof (Sketch):

1137 Part 1: $\exists * \hat{C}$ s.t. $F_{\hat{E}}^\eta(n+1) > F_{\hat{E}}^\eta(n) \forall n \leq \eta \Rightarrow 'x \in R_E' *$ decidable

1138 As before, take any input x and find its new code, $\hat{x}$. By hypothesis the number of steps
 1139 required is bounded by a constant that is independent of system size. Next, we can begin
 1140 to successively evaluate $F_E^\eta(n)$ for increasing values of n . We suppose that the number of
 1141 steps required in this computation for any $n \leq \eta$ is independent of η . This is a reasonable
 1142 assumption because the outputs are identical to those of $\phi_E(n)$ when $n \leq \eta$, and the
 1143 number of steps required to evaluate $\phi_E(n)$ is independent of η for any n . To each output
 1144 of $F_E^\eta(n)$ we can apply the above mapping, $\hat{C}$ to obtain $F_{\hat{E}}^\eta(n)$, which by hypothesis,
 1145 increases with $n \leq \eta$. By hypothesis the number of steps required is independent of η for
 1146 each such application.

1147 As we proceed, either we reach (i) $n = \eta$ prior to reaching an n for which $\hat{x} < F_{\hat{E}}^\eta(n)$, or we
 1148 reach (ii) a value of n whereby $\hat{x} < F_{\hat{E}}^\eta(n)$ before $n = \eta$. In either case $'x \in R_{\hat{E}}'$ is then
 1149 decidable because, if $\hat{x}$ has not been reached by this point, it never will be. Thus, $'x \in R_E'$
 1150 is decidable as well. Moreover, if (i) pertains, then the number of steps required before
 1151 deciding is no more than $\mu i(\phi_{\hat{E}}(i) \geq \hat{x})$. If (ii) pertains, then this number of steps is exactly
 1152 equal to $\mu i(\phi_{\hat{E}}(i) \geq \hat{x})$. And because $\mu i(\phi_{\hat{E}}(i) \geq \hat{x})$ is finite and independent of η , we can
 1153 see that $'x \in R_E'$ is $*$ decidable as well.

1154 Part 2: $'x \in R_E' *$ decidable $\Rightarrow \exists * \hat{C}$ s.t. $F_{\hat{E}}^\eta(n+1) > F_{\hat{E}}^\eta(n) \forall n \leq \eta$

1155 We can construct the required $*$ computable bijection between population states and an
 1156 appropriate coding as follows. First, take any effective coding of population states. By
 1157 hypothesis, the number of steps required to decide $'x \in R_E'$ for any x is finite and

1158 independent of η . Thus, we can proceed through the population states, x , in increasing
1159 order, applying the following algorithm:

1160 (i) if $x \notin R_E$ and it is the k^{th} such state up to that point, use the k^{th} odd number as its
1161 new code.

1162 (ii) if $x \in R_E$, calculate $\mu i(F_E^\eta(i) = x)$, and use the i^{th} even number as its new code.

1163 As we proceed through the states, x , the number of steps required for each, regardless of
1164 whether (i) and (ii) pertains, is independent of η . Therefore, the entire coding procedure
1165 for any given state is independent of η as well; i.e., the coding is *computable as required.